

Bulletin of the Allahabad Mathematical Society

Volume 34, Part 2, 2019

CONTENTS

Saurabh Porwal, Poonam Sharma and Omendra Mishra

HARMONIC MULTIVALENT FUNCTIONS WITH POSITIVE COEFFICIENTS 147-158

Abstract: In the present paper, we introduce and study a new subclass $RS_H^p(n, \lambda, \beta)$ of harmonic multivalent functions by using multiplier transformation. We determine coefficient estimates, extreme points, distortion bounds, convolution properties and convex combination for the above class of harmonic functions and discuss a class preserving integral operator.

Shantanu Bhaumik and Surendra Kumar Tiwari

FIXED POINT THEOREMS FOR MAPPINGS SATISFYING INTEGRAL TYPE
 F -CONTRACTION IN METRIC SPACES ENDOWED WITH A GRAPH 159-172

Abstract: The purpose of this paper is to find the condition for which a function satisfying integral type F - G -contraction defined on a metric space will be Picard operator. With suitable examples we prove that an integral type F - G -contraction is more general than that of an F -contraction and G -contraction.

Mukesh Kumar Singh and Saurabh Porwal

SOME SUFFICIENT CONDITIONS FOR AN INTEGRAL OPERATOR INVOLVING
BESSEL FUNCTIONS 173-180

Abstract: The purpose of the present paper is to obtain some sufficient conditions for starlikeness of an integral operator involving Bessel functions of the first kind. Relevant connections of the results presented here with various known results are briefly indicated.

T. D. Narang and Sahil Gupta

ON CHEBYSHEV CENTERS 181-199

Abstract: In this paper, we survey the main known results on Chebyshev centers and point out some of the connections between the theory of Chebyshev centers and other problems of Approximation Theory and of Banach space theory. We also discuss results on the existence, uniqueness and nature of Chebyshev center points, continuity of the Chebyshev center map and conditions under which bounded sets are centrable.

Om P. Ahuja, S. Sivasubramanian, Y. Polatoğlu and A. Çetinkaya

CERTAIN SUBCLASSES OF BI-UNIVALENT FUNCTIONS GENERATED BY
CHEBYSHEV POLYNOMIALS AND q -DERIVATIVE OPERATOR 201-216

Abstract: In this paper, we introduce and investigate new subclasses of the function class Σ of bi-univalent functions defined in the open unit disc, which are associated with

q -difference operator by means of Chebyshev polynomials. In particular, we find the initial estimates of $|a_2|$ and $|a_3|$ for functions in the new subclasses introduced here. We also obtain Fekete-Szegő inequalities for these classes. Several (known or new) consequences of the results are also obtained.

Sandra Pinelas and Paolo Emilio Ricci

CIRCULAR BASED SHEFFER POLYNOMIAL SEQUENCES

217-234

Abstract: In this article we introduce Sheffer polynomial sequences whose characteristic elements are based on circular or inverse circular function and we derive their main properties, including recurrence relations and differential equations. The article is part of an attempt to sort Sheffer's polynomial sequences on the basis of the characteristics of their generating function. In previous articles, Sheffer sequences arising from other types of generating functions, based on powers or exponentials have been introduced.

Feng-Zhen Zhao

LOG-BEHAVIOR FOR SOME SEQUENCES RELATED TO CAUCHY NUMBERS OF TWO KINDS

235-242

Abstract: In this paper, we discuss log-behavior for some sequences related to Cauchy numbers of two kinds. For Cauchy numbers of two kinds $\{a_n\}_{n \geq 0}$ and $\{b_n\}_{n \geq 0}$, we prove that two sequences $\{|a_n|\}_{n \geq 1}$ and $\{b_n\}_{n \geq 0}$ are 2-log-convex. In addition, we show that sequences $\{|a_{n+1}| - |a_n|\}_{n \geq 2}$, $\{b_{n+1} - b_n\}_{n \geq 1}$ are log-convex, and $\{|a_n|\}_{n \geq 1}$ and $\{b_n\}_{n \geq 0}$ are ratio log-concave.

Eskandar Ameer and Muhammad Arshad

FIXED POINT RESULTS OF GENERALIZED (ψ, ϕ) -RATIONAL TYPE CONTRACTIONS WITH APPLICATION TO NONLINEAR MATRIX EQUATIONS

243-261

Abstract: The purpose of this paper is to introduce the notion of generalized (ψ, ϕ) -rational type contractions and establish some new fixed point theorems for such mappings in complete metric spaces. Our results are an extension and improvement of Jleli and Samet fixed point theorem, Wardowski fixed point theorem, Piri and Kumam fixed point theorem, Geraghty fixed point theorem and Liu et al. fixed point theorem. We give an example for supporting these results. Finally, an application is given to set up the existence of positive definite solution of non-linear matrix equation.
