

CONTENTS

U. C. De and Absos Ali Shaikh

SASAKIAN MAINFOLD WITH C-BOCHNER CURVATURE TENSOR 131-137

Abstract: The object of the present paper is to study sasakian manifolds satisfying the condition $R(X, Y).B = 0$, where B is the C-bochner curvature tensor and $R(X, Y,)$ is considered as a derivation of the tensor algebra at each point of the tangent space. Also sasakian manifolds in which the divergence of the C-bochner curvature tensor vanishes studied.

Julian Dontchev, Maximilian Ganster and Ivan Reilly

MORE ON ALMOST S-CONTINUITY 139-146

Abstract: The aim of the paper is to continue the study of almost s-continuous functions initiated recently by noiri et al.[18]. We improve some of their result and present some new ones. The class of quasi-open sets is introduced.

Server Silvestru Dragomir

ON BEST APPROXIMATION IN REAL NORMED LINEAR SPACES AND CHARACTERIZATIONS OF REFLEXIVITY AND STRICT CONVEXITY 147-158

Abstract: Some characterizations of best approximation element, of proximinal, semitchebychevian and tchebychevian linear subspaces in real normed spaces are given.

H. W. Gould and J. Wetweeraong

EVALUATION OF SOME CLASSES OF BINOMIAL IDENTITIES AND
TWO NEW SETS OF POLYNOMIALS

156-190

Abstract: We prove two main theorems:

1. There exist polynomials $Q_i(n, p)$ of degree $p + 1 - i$ in n such that

$$\sum_{k=j}^n k^p \binom{k}{j} = \sum_{i=1}^{p+1} (-1)^{i-1} Q(n, p) \binom{n+1}{j+i},$$

with $Q_i(n, p) = 0$ for $i < 1$ or for $i > p + 1$. They satisfy the recurrence $Q_{i+1}(n, p) = Q_i(n, p) - Q_i(n - 1, p)$. Also $Q_1(n, p) = n^p$ and $Q_{p+1}(n, p + 1) = p!$.

$$\text{Explicitly } Q_i(n, p) = \sum_{r=0}^{i-1} (-1)^r \binom{i-1}{r} (n - r)^p.$$

2. There exist polynomials $R_i(j, p)$ of degree p in j such that

$$\sum_{k=j}^n \binom{k}{j} = \sum_{i=1}^{p+1} R_i(j, p) \binom{n+1}{j+i}, \text{ with } R_i(j, p) = 0 \text{ for } i < 1.$$

or $i > p + 1$. The R 's satisfy the recurrence $R_i(j, p + 1) = jR_i(j, p) + (j + 1)R_{i-1}(j + 1, p)$. Also $R_1(j, p) = j^p$ and $R_{p+1}(j + 1) = (j + 2) \cdots (j + p) = j!(j + p)_j$

Explicitly

$$R_i(j, p) = \sum_{s=j}^{i+j-1} (-1)^{i+j-s-1} \binom{i+j}{s+1} \sum_{k=j}^s k^p \binom{k}{j}$$

Numerous consequences of these expansions and their connections with the Stirling numbers are presented. Programs using the language C++ have been devised to compute tables of the Q and R polynomials efficiently.

Mona Khare

METRIC ENTROPY AND SUFFICIENT FAMILIES

191-204

Abstract: The aim of the present paper is to introduce the concept of sufficient families in the study of metric entropy $h(\phi, \mathcal{N})$ of

an F-measure preserving transformation ϕ relative to a sub σ -algebra \mathcal{N} of an F-dynamical system $\Phi = (X, M, m, \phi)$, having finitely many atoms. result including the rokhlin inequality have been obtained. It is proved that if \mathcal{N} is a one- sided generator for $[\mathcal{L}]$ with respect to ϕ then the entropy $h(\Phi, [\mathcal{L}]) = 0$. This result includes the corresponding classical result as a particular case.

Byung-Soo Lee and Mee-Kwang Kang

A GENERALIZATION OF SOM AND MUKHERJEE'S FIXED POINT
THEOREM

205-209

Abstract: In this paper a fixed point theorem which generalizes a main result of som and mukherjee [6] is obtained.

Yi-Hai Ma

ON A CLASS OF MIXED MONOTONE OPERATORS AND A KIND OF TWO-POINT
BOUNDARY VALUE PROBLEM

211-220

Abstract: Fixed point theorems for an important class of mixed monotone operators are proved and an application to the two-point boundary value problems for second order impulsive diffrencial equation is given.

S. M. Mazhar

ABSOLUTE SUMMABILITY FACTORS FOR FOURIER SERIES

221-229

Abstract: A theorem on absolute summability factors for fourier series is obtained. Our result generalize certain theorem due to dikshit and dubey and other.

B. Pal, J. C. Misra A. Pal and A. S. Gupta

HYDROMAGNETIC FLOW OF A VISCOELASTIC FLUID IN A PARALLEL PLATE
CHANNEL WITH STRETCHING WALLS 231-247

Abstract: An analysis is made of the study flow of an incompressible viscoelastic and electrically conducting fluid in a parallel plate channel in the presence of a uniform transverse magnetic field. Both the channel walls are stretched in their own plane with a velocity varying linearly with distance from a fixed point. Solution for the velocity distribution is obtained by integrating numerically the ordinary differential equation derived from the governing equations by suitable transformation. For small value of Reynolds number R , the ordinary differential equation also admits of an analytical solution which shows close agreement with the numerical solution. It is found that for fixed R and M (M being the magnetic parameter). The wall shear stress τ_w decreases very slowly with increase in the viscoelastic parameter K_1 . Further it turns out that τ_w increases with increase in M for fixed R and K_1 . A novel result of the present analysis is that the reverse flow arising out of the streamwise velocity changing sign in the transverse direction in the absence of magnetic field ($M = 0$) can be almost eliminated by the application of a strong magnetic field.

G. K. Panda, N. Dash and B. B. Sahoo

ON A GENERALIZED VARIATIONAL INEQUALITY IN HAUSDORFF
TOPOLOGICAL VECTOR SPACES 249-257

Abstract: Recently Singh et al. [6] have given a very beautiful generalization of a result of F.E. Browder on the existence of solution of a variational inequality. Problem using pseudo-monotone mappings in the setting of a Hausdorff topological vector space Behera and Panda [1,2] have also studied several generalizations of Browder's result by considering variational-like inequalities. The aim of this paper is to study one more generalization of a result of Singh et al. which includes some result of Behera and Panda and Siddiqi et al. [5] as cases.

K. P. R. Rao and N. Srinivasa Rao

A COINCIDENCE THEOREM FOR MULTI-VALUED MAPS UNDER
ISHIKAWA ITERATES

259-263

Abstract: In this paper we obtain a common coincidence point theorem for a pair of multi-valued maps with respect to a single-valued map in a normed linear space satisfying a generalized contractive condition using Ishikawa type iteration procedure.

D. Rath

SPACES OF r -CONVEX SEQUENCES AND MATRIX TRANSFORMATIONS 265-280

Abstract: A representation theorem for elements of l_1 in terms of r -convex sequences is obtained which is used to establish matrix characterization theorems generalizing earlier results of Hahn, Dawson. the sequences spaces M_r, bv_p are defined, certain related matrix classes are characterized and the α -duals, β -duals of certain sequences spaces are determined.

B. E. Rhoades

NORMS AND SPECTRA OF GENERALIZED HAUSDORFF MATRICES BOUNDED
ON l^2 AND c

281-306

Abstract: It is shown that certain generalized Hausdorff matrices are bounded operators on l^2 , and a formula is obtained for their l^2 -norm. This paper extends some of the results of [11] to generalized Hausdorff matrices.

Ian Tweddle*

A PERTURBATION THEOREM FOR BARRELLED COUNTABLE ENLARGEMENTS
307-311

Abstract: A barrelled countable enlargement (BCE) is obtained when the dual of a barrelled space is extended by a countably infinite number of dimensions in such a way that barrelledness is preserved. here we consider how the extending vectors of a BCE may be perturbed to produce a new BCE.
